

**Size, age, and species composition of  
commercial eel catches from South Island  
market sampling, 1997–98**

**M. P. Beentjes**



**NIWA Technical Report 51**  
**ISSN 1174-2631**  
**1999**

**Size, age, and species composition of  
commercial eel catches from South Island  
market sampling, 1997–98**

**M. P. Beentjes**

**NIWA Technical Report 50  
1999**

**Published by NIWA  
Wellington  
1999**

Inquiries to:  
Publication Services, NIWA,  
PO Box 14-901, Wellington, New Zealand

**ISSN 1174-2631  
ISBN 0-478-08472-2**

**© NIWA 1999**

Citation: Beentjes, M. P. 1999:  
Size, age, and species composition of commercial eel catches  
from South Island market sampling, 1997–98.  
*NIWA Technical Report 51*. 51 p.

Cover photograph of a female longfinned eel at Lake Wanaka by Michael Beentjes

*The National Institute of Water and Atmospheric Research  
is New Zealand's leading provider  
of atmospheric, marine,  
and freshwater science*

Visit NIWA's website at <http://www.niwa.cri.nz>

# Contents

|  | <i>Page</i> |
|--|-------------|
| Abstract...                                    | 5           |
| Introduction                                   | 5           |
| Methods...                                     | 7           |
| Areas sampled and stratification...            | 7           |
| Sampling procedure                             | 7           |
| Otolith collection and preparation             | 7           |
| Sex...   | 8           |
| Length-weight relationship and condition index | 8           |
| Age and annual growth increments               | 8           |
| Calculation of overall means                   | 9           |
| Results ...                                    | 9           |
| Landings                                       | 9           |
| Length frequency distributions                 | 9           |
| Weight and condition                           | 10          |
| Sex and maturity                               | 11          |
| Age and growth ...                             | 12          |
| Discussion                                     | 13          |
| Landings                                       | 13          |
| Escape tubes                                   | 13          |
| Species composition                            | 13          |
| Size   | 14          |
| Sexual maturity ...                            | 14          |
| Age and growth                                 | 15          |
| Acknowledgments                                | 16          |
| References                                     | 16          |

## Abstract

**Beentjes, M. P. 1999: Size, age, and species composition of commercial eel catches from South Island market sampling, 1997–98.**  
*NIWA Technical Report 51. 51 p.*

This report presents the results of the third consecutive year of market sampling of commercial freshwater eel (shortfinned, *Anguilla australis*; longfinned, *A. dieffenbachii*) landings from throughout the South Island, New Zealand: 104 landings from 11 catchments and 31 strata were sampled and length and weight were recorded for 7869 longfins and 3052 shortfins. The most intensively sampled areas included Te Waihora, and the Maitai, Clutha, and Oreti Rivers. Longfins were the predominant species in all strata except Te Waihora, Lake Brunner, and Waipori Lakes, the proportion usually increasing with distance inland. Longfins were present in 28 and shortfins in 25 of the 31 strata. Length frequency distributions of longfins were generally unimodal with few large eels: shortfin distributions contained a greater proportion of large eels. Overall mean strata length and weight were 55.8 cm and 526 g for longfins and 64.1 cm and 648.7 g for shortfins. Mean size was significantly larger for longfins from less accessible lightly fished areas where females predominated.

Mean age, mean annual weight, and mean annual length increments were determined for eels in two weight categories (220–260 g, 450–550 g). Strata mean ages for longfins in these categories were 20.5 y and 24.5 y respectively, and for shortfins 15.6 y and 17.8 y respectively. The corresponding strata mean annual length and weight increments in these weight categories were 21.7 and 21.8 mm and 12.7 and 16.2 g for longfins, and 28.6 and 34.4 mm and 16.9 and 30.2 g for shortfins. The methods of determining mean age and mean annual weight and length increments from eels within restricted weight categories has been shown to be both cost effective and practical for comparing growth between areas.

The mean strata proportions 35.0, 35.8, and 29.2% of longfins were categorised as immature or unable to determine, male or female; and shortfins 0.1, 4.3, and 95.6% respectively. They indicate that the shortfin fishery, with the exception of Te Waihora, is almost exclusively based on females and the longfin fishery is based mainly on males (ratio of actual numbers of longfin males to females landed was 2.3:1).

Gonad staging indicated that gonad development was related to weight and length and that at maturity, just before migration, longfin males were on average 65 cm, 700 g, and about 23 y; longfin females were 94 cm, 2500 g and 35 y; shortfin males were about 40 cm, under 131 g, and 12–21 y; shortfin females were 82.5 cm, 1100 g, and 30 y.

## Introduction

This report presents the results of the third consecutive year of catch sampling of commercial freshwater eel (shortfinned, *Anguilla australis*; longfinned, *A. dieffenbachii*) landings from throughout the South Island, New Zealand.

The commercial freshwater eel fishery in New Zealand began in the mid 1960s and has been managed largely on the basis of restrictions on permits. Landings peaked in the mid 1970s with the maximum landing of 2434 t recorded in 1975. Landings in recent years averaged about 1400 t (Annala *et al.* 1998) with an export value in 1996 of \$9.5 million. Stock assessment has been limited to interpretation of annual catch data, and the sustainability of the fishery under current levels of harvest is unknown. The general consensus of the eel industry is that the fishery is showing signs of serial depletion in many areas, with fishers exploiting remote and previously unfished areas.

Additionally, despite the absence of quantitative analyses of CPUE, indications are that increasing effort is required to sustain current catch levels. Coupled with this is the widely held view that the size of eels available to both commercial and traditional fishers is steadily declining. As part of the Ministry of Fisheries contract in 1995–96 NIWA examined processors' historical records of eel size grades to address the issue of declining size and changes in species composition (Beentjes & Chisnall 1997). The most comprehensive data sets indicated a clear and progressive trend of declining size from the 1970s through to the present, particularly for longfinned eels. North Island data indicated a decline in the proportion of longfinned eels processed over time, but there was no such indication for the South Island.

Freshwater eels have a high priority for introduction into the Quota Management System and the limited stock assessment information presents problems in determining sustainable harvest levels. Given that processors' historical data indicate a decline in size over time, it is important to monitor size and species composition, together with age and growth of the commercially fished eels within priority catchments. Providing stock assessment advice on a catchment basis is desirable in terms of the independence of stocks and the proposed future management. Growth is a key parameter required for stock assessment in each catchment because it is extremely variable and is highly dependent on factors such as water temperature, food supply, and density (Horn 1996). Ideally, growth should be determined for each catchment (fishery) to provide an index of productivity and to allow changes in growth over time to be monitored.

A pilot programme to monitor the size and species composition of commercial eel stocks within the catchments was implemented in the 1995–96 fishing year in both the South and North Islands. The main goal was to initiate a time-series database on size and species composition and to develop a sampling methodology that could be applied in future years. To determine priority areas, the contribution to eel landings by key areas was determined through a landing survey questionnaire sent to all commercial eel fishers in 1995–96 (Beentjes & Chisnall 1997).

The 1995–96 pilot and 1996–97 programmes sampled 125 and 102 commercial landings respectively, from throughout the South Island. The experience gained from the pilot catch sampling programme in methods used, stratification, and developing relationships with processors and participating fishers was used in the design and operation of the 1996–97 and 1997–98 programmes. Preliminary results of objective 1 to develop an optimal sampling design for the programme were also implemented where possible in 1997–98.

This research was carried out by NIWA under contract to the Ministry of Fisheries as part of Monitoring of eel fisheries (MFish Project No. EEL9701). The programme objective is to assess and monitor commercial eel fisheries and the 1997–98 objective covered in this report was objective 3: 'To monitor the species composition, size structure, and age at the minimum legal size and well above minimum legal size of priority commercial eel fisheries by sampling from 100 landings at the major eel processing shed in the South Island.'

## **Methods**

The 1997–98 South Island catch sampling programme, like those of 1995–96 and 1996–97, was based at Mossburn Enterprises Ltd (Invercargill). The sampling strategy was similar to that used in previous years and was aimed at providing data on eel species, size (length and weight), age, and sex by individual catchment or sub-catchment, some of which were broken into area strata.

### **Areas sampled and stratification**

Selecting which catchments and the number of landings to sample was based on: the 1995–96 landing survey questionnaire results (Beentjes & Chisnall 1997); a need to sample a wide geographical area within the South Island covering habitats representative of both species; availability of cooperative eel fishers who fish from these selected catchments and land into Mossburn Enterprises; the contract requirement to sample 100 landings; preliminary results of the optimal sampling design analysis (objective 1).

Preliminary results on optimal sampling (objective 1) tended to support the practice of dividing catchments into strata. Selected catchments and tributaries were divided into up to four strata based on fishing practices, physical features (e.g., lakes, falls, confluences, weirs, dams, bridges) and information on species and size distribution (Figures 1, 2, Table 1). Stratum boundaries are unchanged from those in 1995–96 and 1996–97 (Beentjes & Chisnall 1997, 1998). Participating fishers were provided with definitions of the boundaries of all strata, Allflex cattle ear tags inscribed with the fisher's name, catchment, and stratum number, and written instructions with details of sampling requirements. Fishers were instructed to keep all eels caught within defined strata separate, and to label these eel samples with cattle ear tags. Eels were delivered by the fisher in holding bags live to the factory at Mossburn Enterprises Ltd, or picked up by the processor's truck which has six aerated tanks; the latter eels were often transported free-swimming. Tags were either tied directly to holding bags or were attached to a float for free-swimming catches in the truck.

### **Sampling procedure**

At the factory, total landed weight (species unsorted) was recorded and a sample taken by randomly selecting several of the holding bags, or by dip netting 100–200 free-swimming eels from the tanker truck. For smaller landings the entire catch was sampled. Eels were deslimed before being processed, with a resultant weight loss estimated at about 3%. Species, length, weight, sex, and maturity were recorded for all eels in the sample. The sample usually contained both species and these were sorted as the sample was analysed. The proportion of each species by weight in the total landed weight was calculated from the proportion by weight of that species in the sample. A record was kept of any eels over 4 kg which were released by fishers: maximum legal size in the South Island is 4 kg (The Fisheries (Commercial Fishing) Regulations, 1986)

### **Otolith collection and preparation**

In 1995–96 and 1996–97, otoliths were collected over the entire size range of eels from each area and this resulted in the collection of up to 150 otoliths per stratum. In 1997–98 the contract objective was to determine age at Minimum Legal Size (MLS, 220 g ) and well above MLS. As a first step the

optimal number of otoliths to sample was determined from the 1996–97 ageing data. A nominal target standard error of under 0.9 was set. The results indicated that a sample of 20 otoliths per size class would satisfy this requirement with a margin of error required for broken and unreadable otoliths. Work on determining the optimum design for catch sampling eels (Francis 1999) indicated that sampling of otoliths should also be spread over as many landings as possible since between landing variance of age was greater than within landing variance of age. Data from 1995–96 and 1996–97 were examined to determine the size range that would be required to achieve a sample size of 20 otoliths from a minimum of three landings per size. The results indicated that the weight categories should be set at 200–240 g and 450–550 g. A South Island industry decision to increase escape tube size from 28 mm to 31 mm in the 1997–98 season meant that the smaller weight category based on MLS had to be increased slightly to 220–260 g.

No attempt was made to determine growth for males and females separately, and the target of 20 otoliths came from one or both sexes. Generally, otoliths were collected from eels in one stratum per catchment where these were mainstem rivers. This was usually stratum 1 (coastal strata), which often yielded the most landings and both species were often better represented. Otoliths were prepared using the crack-and-burn method (Hu & Todd 1981). Otolith halves were mounted in silicone rubber sealant on microscope slides and observed at X10–50 magnification under a stereo-microscope using transmitted light. Hyaline zones or winter rings were counted and age was expressed as years spent in fresh water, ignoring the central area of oceanic larval growth (Jellyman 1979).

## Sex

The sex of eels is difficult to determine externally and often internally, and generally studies have not distinguished between the sexes. Microscopic analysis of gonad tissue in 1995–96 confirmed that assignment of male or female from macroscopic observations was possible. In 1996–97, eels were categorised into four categories: unsexed, immature or unable to determine, male, and female. Gonads were staged as 1 to 4 including the category of immature or unable to determine which showed signs of development but could not be categorised as either male or female (*see* Appendix 1).

## Length-weight relationship and condition index

The length-weight relationship for each species for each stratum (area) was determined from  $\ln W = b(\ln L) + \ln(a)$ , where  $W$  is weight (g) and  $L$  is length (cm). Weight was calculated by setting length equal to 45 cm to provide an index of condition for each stratum. (45 cm approximates length at MLS (220 g) for longfinned eels). No adjustment was made for the estimated 3% weight loss resulting from the desliming process.

## Age and annual growth increments

The mean age within each of the two weight categories (220–260 g and 450–550 g) was calculated and these were termed  $A_{240}$  and  $A_{500}$  respectively. Data from 1995–96 and 1996–97 catch sampling were used to supplement data where sample size was small. Additionally,  $A_{240}$  and  $A_{500}$  were determined for catchments sampled in previous years, but not sampled in 1997–98, by sub-sampling the ageing datasets and selecting ages of eels within the specified weight categories.



Mean annual length and weight increments were derived by dividing the length (minus 50 mm; size at recruitment into fresh water) or weight by age and calculating the mean. Length and weight increments were determined for each weight category. Data from 1995–96 and 1996–97 years catch sampling were also used as described above.

## Calculation of overall means

Overall means for variables such as length and weight were expressed in two ways: the first was the mean of the individual means for each stratum (where N is strata) and is termed *strata mean*; the second is the *all eels mean* calculated without regard to stratification (where N is the total number of eels). These overall means are not always equal due to the weighting effect that sample size can have on the *all eels mean*, i.e., strata that were intensively sampled have a disproportionate effect on, for example, the mean length, whereas when calculating *strata mean*, small or large sample sizes have equal weighting. *All eels means* are shown only in tables and are not referred to in the results.

## Results

### Landings

Between 24 November 1997 and 5 March 1998 104 landings were sampled from 11 catchments (*see* Table 1). Subdivision of these catchments resulted in 31 strata being sampled. Eighteen fishers participated in the programme by providing landing details and/or ensuring that the integrity of catches from designated strata was maintained. The number of landings sampled per stratum varied greatly and was dependent on participating fishers. The most intensively sampled catchments were Te Waihora (Lake Ellesmere), and the Mataura, Clutha, and Oreti Rivers (50 landings). Landed weights of longfins and shortfins totalled 11.2 and 15.5 t respectively, and the overall proportion of the landed weights sampled (sum of sample weights/sum of landed weights per species) was 31.6% for longfins and 8.6% for shortfins. The overall ratio of longfinned to shortfinned eel landed weights was 4:1, (excluding Te Waihora) and about 0.7:1 (including Te Waihora). Length and weight were recorded from 7869 longfins and 3052 shortfins (*see* Table 1) and the mean number of fish sampled per landing for longfins was 87 (N = 90, range = 2–190, s.e. = 4.12) and for shortfins 37 (N = 82, range, 1–152, s.e., 5.13). Otoliths were extracted, prepared and read for 482 longfin and 206 shortfin eels.

Longfins were the predominant species in all strata except those in Te Waihora, Lake Brunner (Grey River, stratum 2), and Waipori Lakes (Taieri River, stratum 3). The proportion of longfins exceeded 84% in all strata except stratum 2 of Waitaki River where it was 74.1%. The proportion of longfins tended to be higher in inland strata, and was frequently 100% (*see* Table 1). Longfins were present in 28 and shortfins in 25 of the 31 strata. Longfinned eels were absent only from Te Waihora strata 1, 3, and 4 with only six longfins caught in stratum 2. Shortfinned eels were absent from inland strata of the Waiau, Oreti, Mataura, Taieri, and Grey Rivers, and from Lake Wakatipu.

### Length frequency distributions

Length frequency distributions of longfins are given by stratum in Figures 3–18. Mean lengths, standard errors, and ranges are given by stratum in Table 2. The strata mean length (28 strata) was 55.8 cm.

From the classification of length frequency distributions for 1996–97 (Beentjes & Chisnall 1998), longfins generally fell into one of three types.

Type 1: Strongly unimodal with mode between 40 and 60 cm centred around 50 cm, with few medium sized or large eels: includes fish from Waiau (strata 1, 2), Aparima, Oreti, Mataura, and Clutha Rivers, Waipori Lakes, and Lake Brunner.

Type 2: An underlying mode of similar size to above is clearly evident, but is skewed to the right with a good representation of medium size eels between 60 and 70 cm: includes fish from Waiau (stratum 3), Makarewa River, Waituna Creek, Mokoreta, Pomahaka, and Waitaki Rivers. Distributions from Waimakariri were similar but with a larger mode centred around 54 cm.

Type 3: No clear modes evident and good representation from all size ranges between 40 and 90 cm: includes fish from Waikaka Stream, upper Taieri River (stratum 4), Lake Wakatipu, and upper Grey River (stratum 3).

Length frequency distributions of shortfins are given by stratum in Figures 19–34. Mean lengths, standard errors, and ranges are given by stratum in Table 3. The strata mean length (25 strata) was 64.1 cm.

The mean length of shortfinned eels was generally greater than that of longfinned eels (Tables 2 and 3) and there was considerably more variability in mean length of shortfinned eels both within and between catchments. Shortfin sample size was small for most strata because of the low proportion of shortfin in landings, with the exception of lower Mataura River (stratum 1) and Waipori Lakes (Taieri River, stratum 3), Lake Brunner (Grey River, stratum 1), and Te Waihora where over 100 eels were sampled. Length frequency distributions of shortfins from these four areas do not have the pronounced unimodal shape of longfins, although modes are apparent between 47 and 63 cm (centred at about 56 cm), and are strongly skewed to the right with all sizes up to about 100 cm represented (Figures 24, 29, 30, and 32 respectively). The distribution of shortfins in stratum 2 of Te Waihora (Kaitorete Spit to Kaituna Lagoon) is similar in shape, but these eels were slightly smaller with the mode centred around 51 cm (Figure 32). The distribution of shortfins caught inside stratum 4 (Concession Area where undersize eels are legally harvested during the shortfin male migration period) is strongly unimodal between 33 and 44 cm (centred at 40 cm); these are migrating males (Figure 32). Because so few shortfinned eels were landed from other areas length frequency distributions cannot be interpreted, but lengths fell within the ranges of the areas described above.

## **Weight and condition**

Mean weight, regression coefficients, and condition indices are given by stratum for longfinned and shortfinned eels (*see* Tables 2 & 3 respectively). The longfin strata mean weight (28 strata) was 526 g and the shortfin strata mean weight (25 strata) was 648 g.

Longfinned eels with the lowest mean weight (where N was over 100) were from lower Aparima River (stratum 1) and the highest mean weight was from Lake Wakatipu (*see* Table 2). The lowest and highest shortfinned eel mean weights (excluding the Te Waihora Concession Area and/or where N was under 100) were from Lake Brunner and Waikaka Stream, respectively (*see* Table 3).

The longfin strata mean condition index was 234 g (28 strata) and the shortfin strata mean condition index was 201 g (25 strata). Longfins with the best condition indices were from inland Waiau River (stratum 3) (244 g) and the poorest were from Lake Wakatipu (197 g). Shortfins with the best condition index were from coastal Waiau River (stratum 1) and Waikaka Stream (238 g for both areas) and the poorest from Te Waihora (stratum 1, 177 g).

## Sex and maturity

Of the 7869 longfins sampled, 98.4% were classified into one of three categories: immature or could not be determined, male, and female. Those that were unclassified were destined for live export and were invariably the larger females. The strata mean proportions of longfins that were categorised as immature or could not be determined, male, or female were 35.0%, 35.8%, and 29.2%, respectively (Table 4). Longfins fell predominantly within the size range 43–65 cm with modes between 52 and 56 cm (Figures 3–17). Female longfins, however, were generally scattered over a larger size range, from about 47 to 97 cm, with no clear modes. The largest eels, although unclassified, were undoubtedly females, which would extend the female upper size range to 110 cm. Eels over 4 kg (equivalent to about 115 cm) must be released by regulation in the South Island, so they were not sampled. The incidence of the category ‘immature or could not be determined’ declines at about the same size that females appear in any number and it is likely that most of these eels will develop into females with immature eels less than 45 cm developing into either male or female.

Although the strata mean proportion of longfin males was 35.8%, some areas had very few males: these include Waikaka Stream (1.1% male, 95.5% female), Lake Wakatipu (0% male, 88.6% female), and upper Grey River (stratum 3; 2.4% male, 89.4% female) (*see* Table 4). The longfin eels from these areas were substantially longer than those from other strata, reflecting the predominance of larger females (*see* Table 2).

As longfinned eels grow in length, gonads mature and change from immature, where the sex cannot be determined, through to the mature stage 4 condition (Table 5). The sex of longfins was clearly distinguishable at mean lengths of about 58 cm for females and 50 cm for males. The stage 4 gonad condition was usually found in both males and females that exhibit morphological signs of migrating, such as enlarged eyes and the shovel-shaped head (Todd 1974, 1980). For South Island longfinned males, the mean length of 64.6 cm at stage 4 corresponds to length at migration. The equivalent length for migrating females is likely to be slightly larger than the mean length of 86.7 cm recorded for stage 4 females because the largest eels were either not sexed, due to processors’ requirements to live export large eels, and/or eels over 4 kg were not landed.

Of the 3052 shortfins sampled, only 2 were not sexed. The remaining eels were classified into three categories: immature or unable to determine, male, and female. The strata mean proportions of shortfins in these categories were 0.1%, 4.3%, and 95.6% respectively (*see* Table 4). Males landed from the Te Waihora Concession Area accounted for 91.7% (N = 603) of all males: the remaining 54 eels came from Te Waihora, and the Mataura, Taieri, and Grey Rivers. There were few immature shortfin males, indicating that shortfin differentiation is occurring at a smaller size than the smallest shortfin males sampled.

Unlike longfins, shortfinned males showed no relationship between gonad maturity and mean length (*see* Table 5). The bulk of the shortfinned males were sampled from the Te Waihora Concession Area where the predominant gonad stage was 3 and all showed morphological signs of migrating described above. Shortfinned males do not grow large and the results on maturity indicate that they migrate

from Te Waihora at a mean length of about 40 cm. Some grading by fishers may occur, so there are probably smaller migrating males in the population than were sampled. Female shortfin eels grow much larger than males and show a clear relationship between gonad maturity and mean length (see Table 5). Length at migration approximates mean length of stage 4 females (82.2 cm) since there were no reports of fishers returning shortfins weighing over 4 kg and only one shortfin (131 cm) was exported live and not sexed.

## Age and growth

Data from 1995–96, 1996–97, and mainly 1997–98 were used to determine longfin mean age and weight within the lower (220–260 g) and upper (450–550 g) weight categories for 19 catchments (not the same 19 catchments in each size range), mostly from coastal strata where these were rivers (Table 6). For the lower weight category 259 otoliths were aged from 42 landings, mean ages ( $A_{240}$ ) ranged from 14.8 y (Waituna Creek) to 30.7 y (Buller River), and the strata mean was 20.5 y. Mean weight ranged from 237.9 to 253.3 g and strata mean weight was 243.5 g. For the upper weight category 283 otoliths were aged from 46 landings, mean ages ( $A_{500}$ ) ranged from 16.9 y (Waitaki River) to 34.8 y (Buller River), and the strata mean was 24.5 y. Mean weight ranged from 478.9 to 540.0 g and strata mean weight was 495 g.

Mean annual length and weight increments for longfins in the two weight categories are given in Table 6. Mean annual length increments ranged from 13.6 mm (Buller River) to 27.9 mm (Waituna Creek) for the lower weight category with a strata mean length increment of 21.7 mm and from 16.2 mm (Buller River) to 31.9 mm (Waitaki River) for the upper weight category with a strata mean length increment of 21.8 mm. Mean annual weight increments ranged from 8.3 g (Buller River) to 16.6 g (Waituna River) for the lower weight category with a strata mean increment of 12.7 g and mean annual weight increments ranged from 16.2 g (Buller River) to 31.9 g (Waitaki River) for the upper weight category with a strata mean increment of 23.1 g. There was little difference in annual length increments between the two weight categories but annual weight increments have generally increased markedly.

Similarly for shortfins, data from 1995–96, 1996–97, and mainly 1997–98 were used to determine shortfin mean age and weight within the lower weight category (220–260 g) for 11 catchments, and the upper weight category (450–550 g) for 14 catchments, mostly from coastal strata where these were rivers (Table 7). For the lower weight category 94 otoliths were aged from 31 landings, mean ages ( $A_{240}$ ) ranged from 10.5 y (Waitaki River) to 20.8 y (Grey River), and the strata mean was 15.6 y. Mean strata weight ranged from 229.2 to 260 g with a strata mean weight of 244.9 g. For the upper weight category 186 otoliths were aged from 85 landings, mean ages ( $A_{500}$ ) ranged from 14.0 y (Oreti and Pomahaka Rivers) to 25.3 y (Grey River), and the strata mean was 17.8 y. Mean weight ranged from 471.7 to 535.0 g with a strata mean weight of 493 g.

Mean annual length and weight increments for shortfins in the two weight categories are given in Table 7. Mean annual length increments ranged from 19.5 mm (Waiau River) to 46.3 mm (Waitaki River) for the lower weight category with a strata mean length increment of 28.6 mm, and from 22.7 mm (Hurunui River) to 50.9 mm (Makarewa River) for the upper weight category with a strata mean length increment of 34.4 mm. Mean annual weight increments ranged from 11.4 g (Waiau River) to 24.2 g (Waitaki River) for the lower weight category with a strata mean increment of 16.9 g and mean

annual weight increments ranged from 19.7 g (Hurunui River) to 37.4 g (Pomahaka River) for the upper weight category with a strata mean increment of 30.2 g. There was a slight increase in annual length increments between the two weight categories but annual weight increments have generally increased markedly.

## Discussion

### Landings

Catch sampling in the South Island was based at Mossburn Enterprises Ltd, Invercargill, where about 65% of all eels landed in the South Island are processed. As in 1995–96 and 1996–97, a high level of cooperation was provided, ensuring that our areal stratification methodology was effective. The total weight of landings sampled in 1997–98 was 4.8% of the average annual South Island commercial landings (Annala *et al.* 1998). Landings sampled in 1997–98 included three new areas: Makarewa River, a tributary of the Oreti River; Waikaka Stream, a tributary of the Mataura River; and Waituna Creek, a small catchment in Southland. The inclusion of these areas, together with other tributaries such as the Mokoreta and Pomahaka Rivers, extends the sampling programme to include less heavily fished areas which tend to be less accessible. Larger eels are generally better represented in these areas (*see* Figure 10) and their inclusion helps to provide a broader view of length distributions and population structure than had sampling been confined to the easily accessed mainstem rivers, the source of most landings. While some catchments from the west coast, Canterbury, Marlborough, and Nelson have been sampled in the last three years (*see* Beentjes & Chisnall 1997, 1998), most landings are from Otago, Southland, and Te Waihora. The last three areas contribute around 70% of the South Island landings (Jellyman 1994).

### Escape tubes

This programme samples only commercially sized or fishery recruited eels. The national MLS of 220 g (excluding Te Waihora) influences the size distribution of eels landed. Catches are seldom effectively manually graded before arriving at the processors. Fishers rely on escape tubes deployed in fyke nets to allow escapement of eels under 220 g. In 1995–96, the minimum legal escape tube diameter was 25 mm, although a code of practice in the South Island encouraged fishers to use 28 mm: this was increased voluntarily to 31 mm in 1997–98. The results from this study indicate that the 31 mm escape tube diameter is effective since only 1.9% and 0.16% (excluding Te Waihora) of longfin and shortfin eels, respectively, weighed less than the MLS of 220 g. In 1996–97, 5% and 0.36% (excluding Te Waihora) of longfin eels and shortfins respectively, weighed less than the MLS of 220 g, indicating that the increase in escape tube diameter from 28 to 31 mm may have contributed to the reduction in numbers of undersized eels landed. Shortfin eels tend to be thinner and weigh less than longfins for a given length, and recruit to the fishery at about 46–47 cm compared to longfins which recruit at around 45 cm.

### Species composition

Longfins were again the predominant commercial species sampled at Mossburn Enterprises, except for landings from Te Waihora, Lake Brunner, (Grey River), and Waipori Lakes which are known shortfin fisheries. Consistent with 1995–96 and 1996–97, proportions of shortfins tended to decline

from coastal to inland strata and sometimes were absent, reflecting habitat preference. Species composition can vary considerably between landings, due in part to the fishing practices of each fisher and the number of landings they contribute to the programme. For example, the proportion of longfins in the lower Mataura River was 83.5% in 1995–96 compared with only 68.1% in 1996–97. One fisher, who tends to target shortfins, contributed few landings in 1995–96 but a significant number in 1996–97 and accordingly the proportion of shortfinned eels from Mataura stratum 1 had increased in 1996–97. In 1997–98 the same fisher contributed only one landing from this stratum and the proportion of longfins was similar to that of 1995–96 (84%). Thus, the apparent annual fluctuations in the proportion of each species from the same area may sometimes be an artefact of sampling.

## Size

South Island length frequency distributions and mean lengths for both species were similar to those in 1995–96 and 1996–97 (*see* Beentjes & Chisnall 1997, 1998). Most longfin distributions are characteristic of exploited fish populations with relatively few large eels. The bulk of the longfin fishery is based on eels between 45 and 60 cm (220–560 g). This is especially evident in the mainstems of southern rivers where length distributions are strongly unimodal with mean lengths around 50 cm. In marked contrast longfin length frequency distributions from less accessible areas which are seldom fished, such as Waikaka Stream, Taieri Gorge, upper Grey River, and Lake Wakatipu, consisted mostly of large females.

Shortfins were, on average, larger than longfins, their length frequency distributions tended not to be unimodal, and catches were more evenly spread over a wider size range. This indicates that fishing pressure has not affected the size structure of shortfin eel populations to the same extent as longfins.

A maximum size limit of 4 kg was introduced in the 1995–96 fishing year for the South Island to protect female longfins. Fishers participating in the catch sampling programme estimated that, from the sampled landings, 114 eels over 4 kg (2 of which were estimated to weigh 10 kg) were caught and released (77 in 1995–96, 116 in 1996–97). The inclusion of these large eels in landings would not have affected the length frequency and age distributions to any extent.

## Sex and maturity

The longfin commercial fishery is dominated by males which are caught 2.3 times as often as females, possibly because females, with their greater longevity, are more vulnerable to fishing. There is also a general decline in the proportion of males from the coast to inland strata, although unlike the American eel, *Anguilla rostrata* (Helfman *et al.* 1987), males were not confined to coastal and estuarine areas. The predominance of females from the upper Waikaka Stream, upper Grey River, and Taieri Gorge may indicate that tributaries and inland areas are the preferred habitat for large females which tend to displace the smaller males.

No longfin males were found in the landing from Lake Wakatipu, where recruitment has been limited since the building of Roxburgh Dam in 1958. Similarly, no males were found in Lake Wakatipu in 1995 (*see* Beentjes *et al.* 1997), but analysis of age indicated that some recruitment occurred after the dam was built. Given the size range of longfinned eels in Lake Wakatipu we would expect to find some males. The total absence of males may support the conclusion that sex determination is

dependent on the environment, and eel populations of low density tend to be largely female while dense populations tend to be predominantly male (*see* Tesch 1977 for review).

Migratory gonad conditions, often associated with morphological changes in migrating eels, indicated that migration in longfins usually takes place at a mean size of about 87 cm (94 cm in 1996–97) and 1.8 kg for females and at 65 cm (65 cm in 1996–97) and 0.7 kg in males, but will vary depending on the fishery. Although these sexed fish were not assigned ages, from the age at length regression plots from 1996–97 (Beentjes & Chisnall 1998), the age of migrating longfins from the lower Maitava River would be about 35 years for females and 23 years for males. These estimates agree well with size and age at migration estimates by Jellyman & Todd (1982), although the female data are slightly biased toward the small size since longfins over 4 kg were returned to the water and not sampled. Additionally, in contrast to 1996–97, some large females were not sexed due to live export requirements and this may account for the apparent difference of 7 cm in length at migration between years.

In contrast to the male dominated longfin fishery, the commercial fishery for shortfins is based almost entirely on females as males migrate at a size below the national MLS (220 g). Some males are still landed from Te Waihora, where the MLS in 1997–98 was 180 g, whereas landings from the Concession Area, where no size limit applies, are almost entirely males. The latter were predominantly migratory males as they displayed clear morphological characteristics of migrating eels and had gonads that were maturing. The mean length and weight were also similar to those of Te Waihora migrating eels recorded in 1993–95 (Jellyman *et al.* 1995). Shortfin gonads differentiate at a smaller size than those of longfins and were distinguishable as female at a mean of 49 cm and male at less than 42 cm, but these males are not vulnerable to fishing except in the Te Waihora Concession Area. The gonad development conditions indicate that female shortfins migrate at a mean length of about 82.2 cm (82.5 cm in 1996–97) and 1.1 kg, and males at 41.7 cm (40.8 cm in 1996–97) and 131 g. Comparable age for females of this length from the Maitava River would be 30 years (Beentjes & Chisnall 1998). The shortfinned migratory males from Te Waihora were aged between 12 and 21 years in 1996–97. These estimates also agree with those of Jellyman & Todd (1982).

## Age and growth

Because growth is highly variable between, and even within, catchments (*see* Beentjes and Chisnall 1998, Francis 1999) ageing was generally confined to the coastal strata of the main rivers with the rationale that ages from these strata should be used as relative indices of growth. Overlaps in the ranges of mean age between the two weight categories are consistent with high variability in age at length or weight. The 1997–98 age data were supplemented with ages from previous years when sample sizes were less than the target sample size of 20. Shortfin eel sample sizes, and particularly the lower weight category, were often less than 20.

From the regression model of age on weight using 1995–96 and 1996–97 data, the mean time to reach MLS (220 g) was 17.5 years for longfins and 12.8 years for shortfins (*see* Beentjes & Chisnall 1998). Because of the increase in escape tube diameter used in 1997–98 the target mean weight of the smaller weight category was increased to 240 g, resulting in mean weights of 243 g for longfin and 245 g for shortfin. The mean time to reach this weight was 20.5 y for longfins and 15.6 y for shortfins; the difference is longer than might be predicted to grow an additional 25 g but may be ascribed to the different statistical techniques used to determine age at a given size. The overall mean time to reach 500 g was 24.5 y for longfins and 17.8 y for shortfins, or about 4.5 years for longfins to grow from 240 to 500 g and just over 2 years for shortfins.

Regression slopes and annual length or weight increments are generally too variable for meaningful comparison and the influence of size on the latter prohibits valid comparisons of productivity between catchments. By restricting the sampling to defined weight categories, annual growth increments between different areas are directly comparable in the same way as mean ages within these categories. Annual length increments were similar for both weight categories (220–260 g, 450–550 g) whereas annual weight increments increased between the small and large weight categories. This reflects the growth characteristic of eels in which the relationship between age and length is often linear but weight increases exponentially in relation to length.

The methods of determining mean age, mean annual weight and length increments from eels within defined size categories have been shown to be both cost effective and practical in terms of comparing growth between areas.

## Acknowledgments

This research was carried out by NIWA under contract to the Ministry of Fisheries (MFish Project No. EEL9701).

Thanks to Victor Thompson and Ron Caughey of Mossburn Enterprises Ltd for allowing us to sample eels in their factory, and to factory staff who cooperated with us. Thanks also to commercial eel fishers who participated in the catch sampling programme by providing landings from defined areas.

We also thank the following NIWA staff: Richard O’Driscoll for coordinating and sampling landings in the South Island, Andrew Willsman for helping collate, analyse, and present the results, and Jill Parkyn for preparing and reading otoliths. Thanks to John Booth for reviewing the report.

## References

- Annala, J. H., Sullivan, K. J., O’Brien, C. J., & Iball, S. D. (Comps.) 1998: Report from the Fishery Assessment Plenary, May 1998: stock assessments and yield estimates. 409 p. (Unpublished report held in NIWA library, Wellington.)
- Beentjes M. P. & Chisnall B. L. 1997: Trends in size and species composition and distribution of commercial eel catches. *New Zealand Fisheries Data Report No. 89*. 71 p.
- Beentjes, M. P. & Chisnall, B. L. 1998: Size, age, and species composition of commercial eel catches from market sampling, 1996–97. *NIWA Technical Report 29*. 124 p.
- Beentjes M. P., Chisnall, B. L., Boubée, J. A. T., & Jellyman, D. J. 1997: Enhancement of the New Zealand eel fishery by elver transfers. *N.Z. Fisheries Technical Report No. 45*. 44 p.
- Francis, R. I. C. C. 1999: Optimum design for shed sampling of eels. New Zealand Fisheries Assessment Research Document 99/3. 28 p. (Draft report held in NIWA library, Wellington.)
- Helfman, G. S., Facey, D. E., Hales, L. S., & Bozeman, E. L. 1987: Reproductive ecology of the American eel. *American Fisheries Society Symposium 1*: 42–56.
- Horn, P. L. 1996: A review of age and growth data for New Zealand freshwater eels (*Anguilla* spp.). New Zealand Fisheries Assessment Research Document 96/6. 23 p. (Draft report held in NIWA library, Wellington.)



- Hu, L. C. & Todd, P. R. 1981: An improved technique for preparing eel otoliths for aging. *New Zealand Journal of Marine and Freshwater Research* 15: 445–446.
- Jellyman, D. J. 1994: The fishery for freshwater eels (*Anguilla* spp.) in New Zealand Fisheries Assessment Research Document 94/14. 24 p. (Draft report held in NIWA library, Wellington.)
- Jellyman, D. J. 1979: Scale development and age determination in New Zealand freshwater eels (*Anguilla* spp.). *New Zealand Journal of Marine and Freshwater Research* 13: 23–30.
- Jellyman, D. J., Chisnall, B. L., & Todd, P. R. 1995: The status of the eel stocks of Lake Ellesmere. *NIWA Science and Technology Series* 26. 62 p.
- Jellyman, D. J. & Todd, P. R. 1982: New Zealand freshwater eels: their biology and fishery. *Fisheries Research Division Information Leaflet* 11. 19 p.
- Tesch, F. W. 1977: The eel. Biology and management of anguillid eels. Chapman and Hall, London. 434 p.
- Todd, P. R. 1974: Studies on the reproductive biology of New Zealand freshwater eels. Unpublished Ph.D. thesis, Victoria University, Wellington. 330 p.
- Todd, P. R. 1980: Size and age of migrating New Zealand freshwater eels (*Anguilla* spp.). *New Zealand Journal of Marine and Freshwater Research* 14: 283–293.

**Table 1 : Catchments and strata sampled in 1997–98 with associated number of samples, landed weights, percent of the landed weight sampled, numbers of eels measured for length and weight, and the proportion of the total landed weight that was longfinned eel (LFE)**

| Catchment      | Stratum definition                        | Stratum | Samples | Longfinned eel |              |                  | Shortfinned eel |              |                  |       |
|----------------|---|---------|---------|----------------|--------------|------------------|-----------------|--------------|------------------|-------|
|                |   |         |         | Landed wt (kg) | % wt sampled | No. eels sampled | Landed wt (kg)  | % wt sampled | No. eels sampled |       |
| Waiau River    | Mouth to Clifden Bridge                   | 1       | 2       | 325.9          | 30.3         | 252              | 12.8            | 40.6         | 11               | 96.2  |
|                | Clifden Bridge to Mararoa Weir            | 2       | 2       | 212.1          | 47.6         | 239              | 16.5            | 55.0         | 14               | 92.8  |
|                | Lake Manapouri to Te Anau control gates   | 3       | 2       | 330.0          | 48.8         | 252              | —               | —            | —                | 100.0 |
| Aparima River  | Mouth to Otautau                          | 1       | 5       | 709.5          | 21.9         | 479              | 54.8            | 40.7         | 31               | 92.8  |
|                | Above Otautau                             | 2       | 4       | 640.4          | 28.1         | 495              | 17.5            | 26.3         | 9                | 97.3  |
| Oreti River    | Mouth to Branxholme                       | 1       | 6       | 446.4          | 55.5         | 570              | 68.3            | 59.7         | 55               | 86.7  |
|                | Branxholme to Centre Bush                 | 2       | 2       | 114.3          | 56.8         | 186              | 1.1             | 36.9         | 1                | 99.0  |
|                | Above Centre Bush                         | 3       | 2       | 445.5          | 21.1         | 278              | —               | —            | —                | 100.0 |
| Makarewa River | Entire river (tributary of Oreti River)   | 1       | 4       | 609.9          | 33.9         | 492              | 33.1            | 28.7         | 16               | 94.9  |
| Waituna Creek  | Entire creek                              | 1       | 1       | 30.6           | 100.0        | 83               | 5.3             | 100.0        | 5                | 85.2  |
| Mataura River  | Mouth to Mataura Falls                    | 1       | 11      | 477.5          | 71.6         | 885              | 91.2            | 77.1         | 117              | 84.0  |
|                | Mataura Falls to Waikata Junction         | 2       | 1       | 95.5           | 47.6         | 97               | —               | —            | —                | 100.0 |
| Mokoreta River | Entire river (tributary of Mataura River) | 1       | 7       | 848.2          | 32.4         | 646              | 57.6            | 34.9         | 36               | 93.6  |
| Waikaka Stream | Entire river (tributary of Mataura River) | 1       | 1       | 295.2          | 29.7         | 94               | 21.2            | 29.7         | 4                | 93.3  |
| Clutha River   | Mouth to Balclutha Bridge                 | 1       | 5       | 945.5          | 20.9         | 571              | 125.1           | 15.6         | 33               | 88.3  |
|                | Balclutha Bridge to Clydevale             | 2       | 5       | 548.2          | 32.3         | 426              | 72.5            | 55.4         | 53               | 88.3  |
|                | Clydevale to Beaumont                     | 3       | 2       | 131.1          | 36.6         | 115              | 5.1             | 29.4         | 2                | 96.3  |
| Pomahaka River | Entire river (tributary of Clutha River)  | 1       | 3       | 446.5          | 28.0         | 221              | 59.9            | 48.4         | 46               | 88.2  |

Table 1 – continued

| Catchment         | Stratum definition                                    | Stratum | Samples | Longfinned eel |              | Shortfinned eel  |                | % landed weight LFE |              |                  |
|-------------------|---|---------|---------|----------------|--------------|------------------|----------------|---------------------|--------------|------------------|
|                   |   |         |         | Landed wt (kg) | % wt sampled | No. eels sampled | Landed wt (kg) |                     | % wt sampled | No. eels sampled |
| Lake Wakatipu     | Headwaters of Clutha River                            | 1       | 1       | 161.0          | 43.4         | 52               | –              | –                   | 100.0        |                  |
| Taieri River      | Waipori Lakes   | 3       | 6       | 124.6          | 60.7         | 185              | 756.2          | 31.9                | 441          | 14.1             |
|                   | Above Taieri Gorge                                    | 4       | 2       | 1012.3         | 13.9         | 218              | –              | –                   | –            | 100.0            |
| Waitaki River     | Mouth to 6 km above Waitaki Bridge                    | 1       | 3       | 424.1          | 34.5         | 237              | 64.7           | 28.9                | 25           | 86.8             |
|                   | 6 km above Waitaki Bridge to Duntroon                 | 2       | 3       | 254.3          | 39.4         | 176              | 89.0           | 46.7                | 66           | 74.1             |
|                   | Duntroon to Waitaki Dam                               | 3       | 1       | 213.4          | 37.7         | 115              | 8.0            | 37.7                | 5            | 96.4             |
| Te Waihora        | Kaituna Lagoon to Halswell River                      | 1       | 1       | –              | –            | –                | 1467.3         | 4.7                 | 128          | 0.0              |
|                   | Kaitorete Spit to Kaituna Lagoon                      | 2       | 5       | 21.6           | 7.9          | 6                | 3591.8         | 7.8                 | 623          | 0.6              |
|                   | Selwyn River to Halswell                              | 3       | 5       | –              | –            | –                | 3359.4         | 6.9                 | 485          | 0.0              |
|                   | Concession Area for migrating males                   | 4       | 5       | –              | –            | –                | 4276.5         | 1.8                 | 626          | 0.0              |
| Waimakariri River | Mouth to Bexley Bridge                                | 1       | 2       | 261.5          | 38.5         | 183              | 1.1            | 28.5                | 1            | 99.6             |
| Grey River        | Lake Brunner and all tributaries that drain into lake | 2       | 4       | 984.2          | 7.8          | 210              | 1239.2         | 7.9                 | 219          | 44.3             |
|                   | Above junction of Arnold and Grey Rivers              | 3       | 1       | 123.1          | 100.0        | 106              | –              | –                   | –            | 100.0            |
| Totals            |   |         | 104     | 11 232.4       |              | 7 869            | 15 495.2       |                     | 3 052        |                  |

**Table 2: Catch sampling length, weight, regression coefficients and condition index for longfinned eels. s.e., standard error; -, insufficient data**

| Catchment      | Stratum | N   | Mean        |      |        | Mean       |       |          | a      | b    | r <sup>2</sup> | Condition index (g) |
|----------------|---------|-----|-------------|------|--------|------------|-------|----------|--------|------|----------------|---------------------|
|                |         |     | length (cm) | s.e. | Range  | weight (g) | s.e.  | Range    |        |      |                |                     |
| Waiau River    | 1       | 252 | 51.7        | 0.53 | 39-99  | 393.2      | 22.11 | 150-3585 | 0.0021 | 3.06 | 0.96           | 221                 |
|                | 2       | 239 | 52.9        | 0.51 | 42-92  | 423.8      | 19.18 | 185-2730 | 0.0019 | 3.08 | 0.94           | 236                 |
|                | 3       | 252 | 58.4        | 0.75 | 44-101 | 645.6      | 33.87 | 225-3725 | 0.0016 | 3.13 | 0.97           | 244                 |
| Aparima River  | 1       | 479 | 50.1        | 0.20 | 42-87  | 327.5      | 5.33  | 190-1840 | 0.0038 | 2.90 | 0.89           | 234                 |
|                | 2       | 495 | 51.5        | 0.25 | 42-102 | 366.6      | 10.40 | 205-3225 | 0.0015 | 3.14 | 0.91           | 228                 |
| Oreti River    | 1       | 570 | 53.4        | 0.32 | 35-109 | 437.6      | 13.77 | 90-3810  | 0.0012 | 3.20 | 0.96           | 231                 |
|                | 2       | 186 | 50.9        | 0.45 | 44-100 | 350.9      | 21.31 | 220-2995 | 0.0015 | 3.12 | 0.91           | 221                 |
|                | 3       | 278 | 50.9        | 0.29 | 42-90  | 347.5      | 8.21  | 215-1825 | 0.0034 | 2.93 | 0.90           | 234                 |
| Makarewa River | 1       | 492 | 52.5        | 0.37 | 40-95  | 425.1      | 13.38 | 200-2775 | 0.0024 | 3.03 | 0.96           | 243                 |
| Waituna Creek  | 1       | 83  | 51.2        | 0.82 | 43-94  | 370.6      | 24.89 | 200-1980 | 0.0056 | 2.80 | 0.93           | 242                 |
| Mataura River  | 1       | 885 | 51.7        | 0.22 | 43-105 | 384.5      | 9.14  | 215-3535 | 0.0019 | 3.08 | 0.93           | 235                 |
|                | 2       | 97  | 55.6        | 0.78 | 45-105 | 478.6      | 35.23 | 250-3420 | 0.0015 | 3.14 | 0.95           | 228                 |
| Mokoreta River | 1       | 646 | 52.6        | 0.32 | 39-100 | 430.6      | 13.03 | 145-3020 | 0.0009 | 3.27 | 0.95           | 232                 |
| Waikaka Stream | 1       | 94  | 68.9        | 1.09 | 49-96  | 934.1      | 52.53 | 290-2955 | 0.0009 | 3.25 | 0.96           | 213                 |
| Clutha River   | 1       | 571 | 50.8        | 0.23 | 41-90  | 349.3      | 6.98  | 195-2055 | 0.0030 | 2.96 | 0.93           | 234                 |
|                | 2       | 426 | 53.4        | 0.35 | 37-98  | 418.4      | 15.20 | 110-3380 | 0.0008 | 3.28 | 0.94           | 218                 |
|                | 3       | 115 | 52.4        | 0.89 | 43-102 | 418.3      | 41.69 | 195-3480 | 0.0009 | 3.26 | 0.97           | 219                 |
| Pomahaka River | 1       | 221 | 57.1        | 0.69 | 42-105 | 570.1      | 33.30 | 190-3705 | 0.0012 | 3.20 | 0.97           | 232                 |

Table 2 – continued

| Catchment         | Stratum | N    | Mean        |      | Range  | Mean weight (g) | s.e.   | Range    | a      | b    | r <sup>2</sup> | Condition index (g) |
|-------------------|---------|------|-------------|------|--------|-----------------|--------|----------|--------|------|----------------|---------------------|
|                   |         |      | length (cm) | s.e. |        |                 |        |          |        |      |                |                     |
| Lake Wakatipu     | 1       | 52   | 72.4        | 2.17 | 50–100 | 1345.3          | 142.10 | 300–3690 | 0.0002 | 3.59 | 0.98           | 197                 |
| Taieri River      | 3       | 185  | 53.2        | 0.50 | 43–110 | 404.9           | 19.37  | 200–3210 | 0.0025 | 3.00 | 0.93           | 230                 |
|                   | 4       | 218  | 61.9        | 0.68 | 47–95  | 640.3           | 26.88  | 250–2570 | 0.0009 | 3.24 | 0.97           | 206                 |
| Waitaki River     | 1       | 237  | 59.9        | 0.64 | 46–101 | 623.8           | 26.36  | 235–3475 | 0.0012 | 3.19 | 0.97           | 226                 |
|                   | 2       | 176  | 59.0        | 0.74 | 44–102 | 571.8           | 32.46  | 205–3570 | 0.0015 | 3.13 | 0.95           | 219                 |
|                   | 3       | 115  | 60.6        | 1.12 | 45–103 | 701.2           | 52.83  | 200–3855 | 0.0008 | 3.30 | 0.98           | 224                 |
| Te Waihora        | 2       | 6    | 49.5        | 2.09 | 44–57  | 287.5           | 39.85  | 195–455  | 0.0023 | 3.00 | 0.92           | 209                 |
| Waimakariri River | 1       | 183  | 57.9        | 0.73 | 45–103 | 554.2           | 31.85  | 205–3515 | 0.0015 | 3.13 | 0.96           | 224                 |
|                   | 2       | 210  | 51.6        | 0.56 | 41–105 | 368.5           | 21.56  | 165–3465 | 0.0013 | 3.15 | 0.95           | 215                 |
| Grey River        | 3       | 106  | 70.2        | 1.38 | 44–99  | 1163.1          | 76.69  | 190–3255 | 0.0005 | 3.40 | 0.97           | 218                 |
| Strata mean       |         |      | 55.8        |      |        | 526.2           |        |          |        |      |                | 234                 |
| All eels mean     |         | 7869 | 53.9        | 0.10 | 35–110 | 454.2           | 4.23   | 90–3855  | 0.0014 | 3.15 | 0.96           | 230                 |

Table 3: Catch sampling length, weight, regression coefficients and condition index for shortfinned eels. s.e., standard error; -, insufficient data

| Catchment      | Stratum | Mean |             |       | Mean   |            |        | a        | b      | r <sup>2</sup> | Condition index (g) |       |
|----------------|---------|------|-------------|-------|--------|------------|--------|----------|--------|----------------|---------------------|-------|
|                |         | N    | length (cm) | s.e.  | Range  | weight (g) | s.e.   |          |        |                |                     | Range |
| Waiau River    | 1       | 11   | 61.1        | 3.09  | 46-81  | 527.7      | 68.16  | 250-990  | 0.0223 | 2.44           | 0.93                | 238   |
|                | 2       | 14   | 65.0        | 2.08  | 50-78  | 667.5      | 60.32  | 295-1155 | 0.0044 | 2.85           | 0.97                | 226   |
| Aparima River  | 1       | 31   | 66.9        | 1.82  | 51-86  | 725.3      | 66.05  | 305-1705 | 0.0012 | 3.14           | 0.96                | 192   |
|                | 2       | 9    | 62.8        | 2.86  | 57-84  | 521.1      | 85.74  | 330-1165 | 0.0016 | 3.05           | 0.98                | 179   |
| Oreti River    | 1       | 55   | 66.9        | 1.48  | 47-91  | 742.1      | 48.03  | 275-1770 | 0.0035 | 2.90           | 0.96                | 217   |
|                | 2       | 1    | -           | -     | 59     | -          | -      | 430      | -      | -              | -                   | -     |
| Makarewa River | 1       | 16   | 63.1        | 2.79  | 49-81  | 602.5      | 76.36  | 285-1160 | 0.0045 | 2.83           | 0.98                | 215   |
| Waituna Creek  | 1       | 5    | 75.2        | 0.94  | 51-104 | 1076.0     | 32.60  | 305-2110 | 0.0064 | 2.74           | 0.98                | 220   |
| Mataura River  | 1       | 117  | 63.5        | 0.94  | 48-97  | 612.5      | 32.59  | 275-2250 | 0.0022 | 3.00           | 0.97                | 200   |
| Mokoreta River | 1       | 36   | 63.5        | 1.28  | 50-79  | 566.5      | 35.21  | 270-1045 | 0.0024 | 2.99           | 0.94                | 210   |
| Waikaka Stream | 1       | 4    | 85.0        | 15.48 | 64-131 | 1570.0     | 799.05 | 565-3955 | 0.0101 | 2.64           | 0.99                | 238   |
| Clutha River   | 1       | 33   | 64.1        | 1.89  | 50-90  | 608.5      | 65.51  | 225-1765 | 0.0010 | 3.18           | 0.98                | 178   |
|                | 2       | 53   | 69.7        | 1.22  | 54-92  | 759.2      | 42.82  | 345-1635 | 0.0026 | 2.95           | 0.95                | 199   |
|                | 3       | 2    | 71.0        | 10.00 | 61-81  | 730.0      | 280.00 | 450-1010 | -      | -              | -                   | -     |
| Pomahaka River | 1       | 46   | 63.5        | 1.74  | 50-95  | 630.2      | 61.46  | 285-2045 | 0.0026 | 2.97           | 0.98                | 205   |
| Taieri River   | 3       | 441  | 62.5        | 0.40  | 47-94  | 551.9      | 12.20  | 200-1820 | 0.0016 | 3.07           | 0.96                | 189   |

Table 3 – continued

| Catchment         | Stratum | N    | length (cm) | s.e. | Range  | weight (g) | s.e.   | Range    | a      | b    | r <sup>2</sup> | Condition index (g) |
|-------------------|---------|------|-------------|------|--------|------------|--------|----------|--------|------|----------------|---------------------|
| Waitaki River     | 1       | 25   | 69.5        | 1.65 | 58–89  | 758.8      | 56.90  | 415–1615 | 0.0048 | 2.81 | 0.90           | 215                 |
|                   | 2       | 66   | 65.2        | 1.15 | 48–97  | 634.9      | 35.71  | 270–1875 | 0.0032 | 2.91 | 0.95           | 204                 |
|                   | 3       | 5    | 62.2        | 3.97 | 53–75  | 596.0      | 131.12 | 330–1080 | 0.0012 | 3.16 | 0.95           | 202                 |
| Te Waihora        | 1       | 128  | 59.9        | 0.85 | 45–86  | 539.9      | 29.77  | 205–1665 | 0.0003 | 3.49 | 0.98           | 177                 |
|                   | 2       | 623  | 55.9        | 0.44 | 31–96  | 453.1      | 13.32  | 50–2285  | 0.0005 | 3.39 | 0.97           | 185                 |
|                   | 3       | 485  | 58.5        | 0.36 | 42–91  | 481.4      | 11.52  | 165–1850 | 0.0008 | 3.25 | 0.96           | 190                 |
|                   | 4       | 626  | 39.5        | 0.11 | 32–56  | 116.1      | 1.32   | 60–365   | 0.0008 | 3.22 | 0.78           | 172                 |
| Waimakariri River | 1       | 1    | –           | –    | –      | –          | –      | 345      | –      | –    | –              | –                   |
| Grey River        | 2       | 219  | 59.2        | 0.51 | 43–85  | 448.6      | 13.35  | 195–1380 | 0.0014 | 3.10 | 0.95           | 182                 |
| Strata mean       |         |      | 64.1        |      |        | 648.7      |        |          |        |      |                | 201                 |
| All eels mean     |         | 3052 | 55.9        | 0.22 | 31–131 | 442.6      | 5.91   | 50–3955  | 0.0006 | 3.31 | 0.98           | 180                 |

**Table 4: Percentage of longfinned and shortfinned eels in each stratum that were male (M), female (F), or immature or unable to determine (I). –, no data**

| Catchment         | Stratum | Longfinned eels |      |      |       | Shortfinned eels |       |     |       |
|-------------------|---------|-----------------|------|------|-------|------------------|-------|-----|-------|
|                   |         | M               | F    | I    | Total | M                | F     | I   | Total |
| Waiau River       | 1       | 47.8            | 11.6 | 40.6 | 249   | 0.0              | 100.0 | 0.0 | 11    |
|                   | 2       | 50.9            | 15.0 | 34.2 | 234   | 0.0              | 100.0 | 0.0 | 14    |
|                   | 3       | 43.8            | 30.8 | 25.4 | 240   | –                | –     | –   | 0     |
| Aparima River     | 1       | 55.3            | 4.8  | 39.9 | 479   | 0.0              | 100.0 | 0.0 | 31    |
|                   | 2       | 52.0            | 4.3  | 43.7 | 492   | 0.0              | 100.0 | 0.0 | 9     |
| Oreti River       | 1       | 49.1            | 18.2 | 32.7 | 566   | 0.0              | 100.0 | 0.0 | 54    |
|                   | 2       | 50.0            | 2.2  | 47.8 | 184   | 0.0              | 100.0 | 0.0 | 1     |
|                   | 3       | 47.8            | 5.4  | 46.8 | 278   | –                | –     | –   | 0     |
| Makarewa River    | 1       | 41.6            | 19.5 | 38.9 | 488   | 0.0              | 100.0 | 0.0 | 16    |
| Waituna Creek     | 1       | 9.6             | 18.1 | 72.3 | 83    | 0.0              | 100.0 | 0.0 | 5     |
| Mataura River     | 1       | 52.6            | 2.4  | 45.0 | 878   | 1.7              | 98.3  | 0.0 | 117   |
|                   | 2       | 39.6            | 31.3 | 29.2 | 96    | –                | –     | –   | 0     |
| Mokoreta River    | 1       | 42.1            | 17.8 | 40.1 | 636   | 0.0              | 100.0 | 0.0 | 36    |
| Waikaka Stream    | 1       | 1.1             | 95.5 | 3.4  | 89    | 0.0              | 100.0 | 0.0 | 3     |
| Clutha River      | 1       | 49.6            | 9.5  | 40.9 | 570   | 0.0              | 100.0 | 0.0 | 33    |
|                   | 2       | 49.4            | 13.1 | 37.5 | 421   | 0.0              | 100.0 | 0.0 | 53    |
|                   | 3       | 30.1            | 15.0 | 54.9 | 113   | 0.0              | 100.0 | 0.0 | 2     |
| Pomahaka River    | 1       | 46.3            | 32.2 | 21.5 | 214   | 0.0              | 100.0 | 0.0 | 46    |
| Lake Wakatipu     | 1       | 0.0             | 88.6 | 11.4 | 35    | –                | –     | –   | 0     |
| Taieri River      | 3       | 46.2            | 16.3 | 37.5 | 184   | 0.7              | 99.3  | 0.0 | 441   |
|                   | 4       | 15.3            | 65.1 | 19.5 | 215   | –                | –     | –   | 0     |
| Waitaki River     | 1       | 31.8            | 43.2 | 25.0 | 236   | 0.0              | 100.0 | 0.0 | 25    |
|                   | 2       | 40.5            | 38.2 | 21.4 | 173   | 0.0              | 100.0 | 0.0 | 66    |
|                   | 3       | 27.3            | 42.7 | 30.0 | 110   | 0.0              | 100.0 | 0.0 | 5     |
| Te Waihora        | 1       | –               | –    | –    | 0     | 0.0              | 100.0 | 0.0 | 128   |
|                   | 2       | 16.7            | 33.3 | 50.0 | 6     | 7.2              | 91.3  | 1.4 | 623   |
|                   | 3       | –               | –    | –    | 0     | 0.2              | 99.6  | 0.2 | 485   |
|                   | 4       | –               | –    | –    | 0     | 96.3             | 3.0   | 0.6 | 626   |
| Waimakariri River | 1       | 36.7            | 37.8 | 25.6 | 180   | 0.0              | 100.0 | 0.0 | 1     |
| Grey River        | 2       | 27.3            | 15.3 | 57.4 | 209   | 1.4              | 98.6  | 0.0 | 219   |
|                   | 3       | 2.4             | 89.4 | 8.2  | 85    | –                | –     | –   | 0     |
| Strata mean       |         | 35.8            | 29.2 | 35.0 |       | 4.3              | 95.6  | 0.1 |       |
| All eels mean     |         | 43.8            | 18.8 | 37.5 | 7743  | 21.5             | 78.0  | 0.5 | 3050  |



**Table 5: Mean length of longfinned and shortfinned eels by gonad stage. See Appendix 1 for gonad descriptions. s.e., standard error**

| Gonad stage | Indeterminate sex |                  |      | Male |                  | Female |                  |
|-------------|-------------------|------------------|------|------|------------------|--------|------------------|
|             | N                 | Mean length (cm) | s.e. | N    | Mean length (cm) | N      | Mean length (cm) |
| Longfins    | 2901              | 48.67            | 0.06 | —    | —                | —      | —                |
| Immature    | —                 | —                | —    | 2463 | 50.3             | 1063   | 58.4             |
| 1           | —                 | —                | —    | 779  | 55.4             | 325    | 72.9             |
| 2           | —                 | —                | —    | 125  | 59.8             | 52     | 81.8             |
| 3           | —                 | —                | —    | 23   | 64.6             | 12     | 86.7             |
| 4           | —                 | —                | —    | —    | —                | —      | —                |
| Totals      | 2901              | —                | —    | 3390 | —                | 1452   | —                |
| Shortfins   | 14.0              | 38.6             | 1.50 | —    | —                | —      | —                |
| Immature    | —                 | —                | —    | 6    | 41.7             | 320    | 48.9             |
| 1           | —                 | —                | —    | 58   | 38.8             | 1875   | 60.2             |
| 2           | —                 | —                | —    | 511  | 38.9             | 168    | 77.4             |
| 3           | —                 | —                | —    | 82   | 39.3             | 16     | 82.2             |
| 4           | —                 | —                | —    | —    | —                | —      | —                |
| Totals      | 14                | —                | —    | 657  | —                | 2379   | —                |

**Table 6: Longfinned eel mean age (A<sub>240</sub>, A<sub>500</sub>) and mean annual length and weight increments, for two weight categories, 220–260 g and 450–550 g. All data were collected in 1997–98 unless otherwise indicated. s.e., standard error; \*, 1997–98 data supplemented from 1995–96, 1996–97; \*\*, data from 1995–96, 1996–97**

| Catchment      | Stratum | Number of landings sampled | N     | Mean weight (g) | Mean length (cm) | Age range | Weight category (g)        | Mean age ± s.e.            |                            | Mean annual weight increment (g) ± s.e. |           | Mean annual length increment (mm) ± s.e. |           |
|----------------|---------|----------------------------|-------|-----------------|------------------|-----------|----------------------------|----------------------------|----------------------------|---|-----------|--|-----------|
|                |         |                            |       |                 |                  |           |                            | A <sub>240</sub>           | A <sub>500</sub>           | 220–260                                 | 450–550   | 220–260                                  | 450–550   |
| Waiau River    | 1       | 1                          | 20    | 243.3           | 45.7             | 18–30     | 220–260                    | A <sub>240</sub> 21.6±1.77 | A <sub>500</sub> 26.8±0.99 | 11.6±0.43                               | 18.4±0.64 | 19.3±0.67                                | 19.8±0.64 |
|                | 2       | 2                          | 12    | 486.7           | 57.5             | 21–33     | 450–550                    |                            |                            |   |           |  |           |
| Aparima River  | 1       | 3                          | 30    | 246.0           | 46.6             | 15–24     | 220–260                    | A <sub>240</sub> 19.1±0.43 |                            | 13.1±0.30                               |           | 22.1±0.48                                |           |
|                |         | 5                          | 18    | 491.3           | 55.0             | 15–35     | 450–550                    | A <sub>500</sub> 25.4±1.28 |                            | 20.3±1.25                               |           | 21.6±1.25                                |           |
|                | 2       | 1                          | 8     | 494.4           | 57.4             | 21–29     | 450–550                    | A <sub>500</sub> 24.1±1.04 |                            | 20.7±0.64                               |           | 21.0±0.78                                |           |
|                |         | 4                          | 22    | 243.4           | 46.2             | 11–23     | 220–260                    | A <sub>240</sub> 18.6±0.55 |                            | 13.4±0.49                               |           | 22.6±0.79                                |           |
| Oreti River    | 3       | 24                         | 494.8 | 56.6            | 17–33            | 450–550   | A <sub>500</sub> 22.9±0.91 |                            | 22.3±0.77                  |   | 23.3±0.90 |  |           |
|                | 1       | 2                          | 243.4 | 45.1            | 12–20            | 220–260   | A <sub>240</sub> 15.8±0.52 |                            | 15.8±0.61                  |   | 26.1±0.93 |  |           |
| Makarewa River | 3       | 21                         | 487.4 | 55.6            | 16–31            | 450–550   | A <sub>500</sub> 21.1±0.83 |                            | 23.8±0.91                  |   | 24.8±0.91 |  |           |
|                | 1       | 1                          | 9     | 245.6           | 46.0             | 13–16     | 220–260                    | A <sub>240</sub> 14.8±0.36 |                            | 16.6±0.45                               |           | 27.9±0.70                                |           |
| Waituna Creek  | 1       | 1                          | 8     | 481.3           | 57.4             | 17–24     | 450–550                    | A <sub>500</sub> 19.9±0.88 |                            | 24.4±0.84                               |           | 26.6±0.87                                |           |
|                | 1       | 6                          | 26    | 252.9           | 46.7             | 13–24     | 220–260                    | A <sub>240</sub> 17.7±0.56 |                            | 14.7±0.44                               |           | 24.2±0.72                                |           |
| Mataura River  | 6       | 22                         | 495.2 | 56.6            | 15–36            | 450–550   | A <sub>500</sub> 21.9±1.28 |                            | 23.8±1.10                  |   | 24.9±1.20 |  |           |
|                | 1       | 4                          | 243.8 | 45.9            | 19–34            | 220–260   | A <sub>240</sub> 26.0±0.88 |                            | 9.6±0.32                   |   | 16.0±0.55 |  |           |
| Mokoreta River | 3       | 20                         | 494.5 | 56.4            | 25–42            | 450–550   | A <sub>500</sub> 32.0±1.08 |                            | 15.8±0.55                  |   | 16.4±0.53 |  |           |
|                | 1       | 1                          | 13    | 509.2           | 59.2             | 14–31     | 450–550                    | A <sub>500</sub> 22.2±1.40 |                            | 24.1±1.73                               |           | 25.6±1.72                                |           |
| Clutha River   | 1       | 2                          | 246.8 | 46.3            | 13–22            | 220–260   | A <sub>240</sub> 17.4±0.48 |                            | 14.3±0.33                  |   | 24.0±0.62 |  |           |
|                |         | 3                          | 20    | 494.0           | 57.4             | 15–32     | 450–550                    | A <sub>500</sub> 23.2±1.06 |                            | 22.3±1.13                               |           | 23.5±1.10                                |           |
|                | 2       | 1                          | 14    | 245.0           | 47.0             | 12–23     | 220–260                    | A <sub>240</sub> 18.9±0.73 |                            | 13.3±0.66                               |           | 22.9±1.19                                |           |
|                |         | 1                          | 10    | 504.4           | 58.5             | 21–34     | 450–550                    | A <sub>500</sub> 25.6±1.12 |                            | 19.9±0.57                               |           | 21.2±0.81                                |           |
| Pomahaka River | 1       | 2                          | 237.9 | 46.3            | 14–21            | 220–260   | A <sub>240</sub> 18.7±1.11 |                            | 13.0±0.87                  |   | 22.6±1.54 |  |           |
|                |         | 3                          | 20    | 484.5           | 57.4             | 15–38     | 450–550                    | A <sub>500</sub> 24.6±1.42 |                            | 21.1±1.33                               |           | 22.8±1.39                                |           |
|                | 1       | 1                          | 7     | 237.9           | 46.3             | 14–21     | 220–260                    | A <sub>240</sub> 18.7±1.11 |                            | 13.0±0.87                               |           | 22.6±1.54                                |           |

Table 6 – continued


| Catchment         | Stratum | Number of landings sampled | N  | Mean weight (g) | Mean length (cm) | Age range | Weight category (g) | Mean age $\pm$ s.e.              | Mean                                   |   |
|-------------------|---------|----------------------------|----|-----------------|------------------|-----------|---------------------|----------------------------------|--|---|
|                   |         |                            |    |                 |                  |           |                     |                                  | annual weight increment (g) $\pm$ s.e. | annual length increment (mm) $\pm$ s.e. |
| Taieri River      | 3       | 1 **                       | 1  | 235.0           | 45.0             | –         | 220–260             | A <sub>240</sub> 19.0            | 14.1 $\pm$ 1.70                        | 25.9 $\pm$ 4.86                         |
|                   |         | 1 **                       | 1  | 500.0           | 57.0             | –         | 450–550             | A <sub>500</sub> 20.0            | 25.0                                   | 26.0                                    |
| Lake Wakatipu     | 1       | 1                          | 5  | 485.0           | 57.0             | 18–28     | 450–550             | A <sub>500</sub> 21.2 $\pm$ 1.93 | 23.4 $\pm$ 1.39                        | 25.1 $\pm$ 1.77                         |
| Waitaki River     | 1       | 4 *                        | 8  | 243.8           | 46.1             | 14–27     | 220–260             | A <sub>240</sub> 17.3 $\pm$ 1.58 | 14.8 $\pm$ 1.07                        | 24.9 $\pm$ 1.77                         |
|                   |         | 3                          | 20 | 500.0           | 57.7             | 13–22     | 450–550             | A <sub>500</sub> 16.9 $\pm$ 0.58 | 30.3 $\pm$ 1.05                        | 31.9 $\pm$ 1.14                         |
| Rangitata River   | 1       | 1 **                       | 4  | 236.3           | 48.3             | 15–20     | 220–260             | A <sub>240</sub> 18.3 $\pm$ 1.11 | 13.2 $\pm$ 1.22                        | 24.1 $\pm$ 2.03                         |
| Rakaia River      | 1       | 1 **                       | 1  | 240.0           | 48.0             | –         | 220–260             | A <sub>240</sub> 18.0            | 13.3                                   | 23.9                                    |
|                   |         | 1 **                       | 5  | 491.0           | 60.2             | 16–22     | 450–550             | A <sub>500</sub> 19.6 $\pm$ 1.29 | 25.5 $\pm$ 1.83                        | 28.6 $\pm$ 1.78                         |
| Te Waihora        | 2       | 1                          | 1  | 540.0           | 61.0             | –         | 450–550             | A <sub>500</sub> 19.0            | 28.4                                   | 29.5                                    |
| Waimakariri River | 1       | 2                          | 5  | 243.0           | 46.8             | 17–34     | 220–260             | A <sub>240</sub> 26.4 $\pm$ 3.17 | 9.8 $\pm$ 1.37                         | 17.0 $\pm$ 2.57                         |
|                   |         | 2                          | 22 | 501.3           | 57.4             | 16–45     | 450–550             | A <sub>500</sub> 29.8 $\pm$ 1.54 | 17.9 $\pm$ 1.11                        | 18.7 $\pm$ 1.13                         |
| Huronui River     | 2       | 1 **                       | 3  | 243.3           | 47.0             | 19–24     | 220–260             | A <sub>240</sub> 21.0 $\pm$ 1.53 | 11.8 $\pm$ 1.19                        | 20.2 $\pm$ 1.65                         |
|                   |         | 1 **                       | 9  | 501.1           | 57.3             | 20–28     | 450–550             | A <sub>500</sub> 25.0 $\pm$ 0.80 | 20.2 $\pm$ 0.79                        | 21.1 $\pm$ 0.68                         |
| Awatere River     | 1       | 1 **                       | 9  | 235.6           | 45.8             | 12–21     | 220–260             | A <sub>240</sub> 17.2 $\pm$ 0.92 | 13.9 $\pm$ 0.74                        | 24.3 $\pm$ 1.41                         |
| Buller River      | 1       | 2 **                       | 3  | 253.3           | 46.3             | 28–35     | 220–260             | A <sub>240</sub> 30.7 $\pm$ 2.19 | 8.3 $\pm$ 0.53                         | 13.6 $\pm$ 0.79                         |
|                   |         | 2 **                       | 11 | 480.0           | 59.5             | 21–54     | 450–550             | A <sub>500</sub> 34.8 $\pm$ 2.68 | 14.5 $\pm$ 1.05                        | 16.2 $\pm$ 0.99                         |
| Grey River        | 2       | 3                          | 24 | 240.4           | 46.4             | 17–32     | 220–260             | A <sub>240</sub> 25.9 $\pm$ 0.71 | 9.4 $\pm$ 0.28                         | 16.3 $\pm$ 0.53                         |
|                   |         | 4                          | 14 | 478.9           | 58.1             | 21–40     | 450–550             | A <sub>500</sub> 33.2 $\pm$ 1.25 | 14.7 $\pm$ 0.68                        | 16.4 $\pm$ 0.85                         |
| Hokitika River    | 1       | 1 **                       | 12 | 242.5           | 47.6             | 20–33     | 220–260             | A <sub>240</sub> 26.2 $\pm$ 1.03 | 9.3 $\pm$ 0.34                         | 16.5 $\pm$ 0.58                         |

**Table 7: Shortfinned eel mean age ( $A_{240}$ ,  $A_{500}$ ) and mean annual length and weight increments, for two weight categories, 220–260 g and 450–550 g. All data were collected in 1997–98 unless otherwise indicated. s.e., standard error; \*, 1997–98 data supplemented from 1995–96, 1996–97; \*\*, data from 1995–96, 1996**


| Catchment      | Stratum | Number of landings sampled | N  | Mean weight (g) | Mean length (cm) | Age range | Weight category (g) | Mean age $\pm$ s.e.       | Mean                                   |   |
|----------------|---------|----------------------------|----|-----------------|------------------|-----------|---------------------|---------------------------|--|---|
|                |         |                            |    |                 |                  |           |                     |                           | annual weight increment (g) $\pm$ s.e. | annual length increment (mm) $\pm$ s.e. |
| Waiau River    | 1       | 1                          | 1  | 250.0           | 48.0             | –         | 220–260             | $A_{240}$ 22.0            | 11.4                                   | 19.5                                    |
|                |         | 3 *                        | 5  | 535.0           | 63.4             | 17–26     | 450–550             | $A_{500}$ 20.4 $\pm$ 1.54 | 26.7 $\pm$ 1.69                        | 29.2 $\pm$ 1.97                         |
| Aparima River  | 1       | 1 **                       | 2  | 260.0           | 49.0             | 14–18     | 220–260             | $A_{240}$ 16.0 $\pm$ 2.00 | 16.5 $\pm$ 2.06                        | 27.9 $\pm$ 3.49                         |
|                |         | 1                          | 3  | 471.7           | 60.0             | 19–23     | 450–550             | $A_{500}$ 20.7 $\pm$ 1.20 | 22.9 $\pm$ 1.34                        | 26.8 $\pm$ 1.95                         |
| Oreti River    | 1       | 1 **                       | 1  | 245.0           | 50.0             | –         | 220–260             | $A_{240}$ 17.0            | 14.4                                   | 26.5                                    |
|                |         | 5 *                        | 10 | 485.0           | 60.5             | 10–18     | 450–550             | $A_{500}$ 14.0 $\pm$ 0.83 | 35.7 $\pm$ 2.11                        | 40.9 $\pm$ 2.58                         |
| Makarewa River | 1       | 1                          | 1  | 485.0           | 61.0             | –         | 450–550             | $A_{500}$ 11.0            | 44.1                                   | 50.9                                    |
| Waituna Creek  | 1       | 1                          | 1  | 475.0           | 57.0             | –         | 450–550             | $A_{500}$ 16.0            | 29.7                                   | 32.5                                    |
| Mataura River  | 1       | 11 *                       | 11 | 246.4           | 49.1             | 8–19      | 220–260             | $A_{240}$ 12.5 $\pm$ 1.04 | 21.0 $\pm$ 1.58                        | 37.8 $\pm$ 3.02                         |
|                |         | 7 **                       | 38 | 492.9           | 60.8             | 8–22      | 450–550             | $A_{500}$ 15.2 $\pm$ 0.65 | 35.0 $\pm$ 1.66                        | 39.8 $\pm$ 1.90                         |
| Mokoreta River | 1       | 2                          | 4  | 495.0           | 61.3             | 22–26     | 450–550             | $A_{500}$ 24.3 $\pm$ 0.85 | 20.5 $\pm$ 1.25                        | 23.3 $\pm$ 1.18                         |
| Clutha River   | 1       | 2 *                        | 2  | 242.5           | 49.0             | 10–14     | 220–260             | $A_{240}$ 12.0 $\pm$ 2.00 | 21.0 $\pm$ 4.96                        | 37.6 $\pm$ 5.43                         |
|                |         | 2                          | 6  | 483.3           | 61.5             | 11–24     | 450–550             | $A_{500}$ 14.7 $\pm$ 1.93 | 34.7 $\pm$ 2.77                        | 40.8 $\pm$ 3.57                         |
| Pomahaka River | 1       | 1 **                       | 1  | 260.0           | 50.0             | –         | 220–260             | $A_{240}$ 16.0            | 16.3                                   | 28.1                                    |
|                |         | 2 *                        | 6  | 474.2           | 60.5             | 8–19      | 450–550             | $A_{500}$ 14.0 $\pm$ 1.79 | 37.4 $\pm$ 5.65                        | 43.5 $\pm$ 6.11                         |
| Taieri River   | 3       | 2 *                        | 3  | 255.0           | 47.7             | 10–16     | 220–260             | $A_{240}$ 13.7 $\pm$ 1.86 | 19.4 $\pm$ 2.82                        | 32.5 $\pm$ 4.82                         |
|                |         | 5 *                        | 27 | 494.3           | 62.1             | 11–33     | 450–550             | $A_{500}$ 15.8 $\pm$ 0.96 | 33.2 $\pm$ 1.33                        | 38.5 $\pm$ 1.58                         |

*Table 7 – continued*

| Catchment     | Stratum | Number of landings sampled | N  | Mean weight (g) | Mean length (cm) | Age range | Weight category (g) | Mean age $\pm$ s.e.              | Mean                                   |   |
|---------------|---------|----------------------------|----|-----------------|------------------|-----------|---------------------|----------------------------------|--|---|
|               |         |                            |    |                 |                  |           |                     |                                  | annual weight increment (g) $\pm$ s.e. | annual length increment (mm) $\pm$ s.e. |
| Waitaki River | 1       | 3 **                       | 6  | 229.2           | 48.7             | 7–19      | 220–260             | A <sub>240</sub> 10.5 $\pm$ 1.80 | 24.2 $\pm$ 2.86                        | 46.3 $\pm$ 5.64                         |
|               |         | 6 *                        | 12 | 505.8           | 62.2             | 10–35     | 450–550             | A <sub>500</sub> 17.5 $\pm$ 1.82 | 31.5 $\pm$ 2.52                        | 35.5 $\pm$ 2.76                         |
| Te Waihora    | 2       | 3                          | 20 | 236.8           | 48.8             | 9–20      | 220–260             | A <sub>240</sub> 14.9 $\pm$ 0.61 | 16.5 $\pm$ 0.74                        | 30.6 $\pm$ 1.44                         |
|               |         | 3                          | 20 | 507.5           | 59.9             | 10–20     | 450–550             | A <sub>500</sub> 16.6 $\pm$ 0.55 | 31.2 $\pm$ 1.13                        | 33.9 $\pm$ 1.33                         |
|               | 3       | 3                          | 20 | 245.0           | 48.5             | 11–20     | 220–260             | A <sub>240</sub> 14.9 $\pm$ 0.46 | 16.8 $\pm$ 0.57                        | 29.8 $\pm$ 0.99                         |
|               |         | 2                          | 21 | 495.7           | 60.1             | 11–20     | 450–550             | A <sub>500</sub> 16.4 $\pm$ 0.58 | 31.1 $\pm$ 1.33                        | 34.5 $\pm$ 1.44                         |
| Huruni River  | 2       | 2 **                       | 6  | 494.2           | 61.8             | 22–30     | 450–550             | A <sub>500</sub> 25.3 $\pm$ 1.15 | 19.7 $\pm$ 0.91                        | 22.7 $\pm$ 1.03                         |
| Wairau River  | 2       | 1 **                       | 7  | 242.9           | 49.7             | 13–22     | 220–260             | A <sub>240</sub> 17.4 $\pm$ 1.23 | 14.4 $\pm$ 1.19                        | 26.5 $\pm$ 1.99                         |
| Grey River    | 2       | 3                          | 21 | 241.4           | 49.3             | 16–28     | 220–260             | A <sub>240</sub> 20.8 $\pm$ 0.66 | 11.9 $\pm$ 0.42                        | 21.8 $\pm$ 0.71                         |
|               |         | 3                          | 26 | 497.7           | 62.4             | 16–35     | 450–550             | A <sub>500</sub> 25.3 $\pm$ 0.89 | 20.3 $\pm$ 0.74                        | 23.5 $\pm$ 0.90                         |



**Figure 1:** Catchments where commercial landings were sampled as part of the South Island catch sampling programme in 1997–98. Strata within catchments are indicated by dotted lines and numerals (see Table 1 for strata definitions).



**Figure 2:** Te Waihora catchment (enlarged from Figure 1) showing strata boundaries used in the South Island catch sampling programme in 1997–98.






Figure 3: Length frequencies of longfinned eels from strata 1, 2, and 3 of Waiiau River.











Figure 4: Length frequencies of longfinned eels from strata 1, 2, and 3 of Oreti River.



**Figure 5: Length frequencies of longfinned eels from stratum 1 of Makarewa River.**



**Figure 6: Length frequencies of longfinned eels from strata 1 and 2 of Aparima River.**




Figure 7: Length frequencies of longfinned eels from stratum 1 of Waituna Creek.





Figure 8: Length frequencies of longfinned eels from strata 1 and 2 of Matura River.




Figure 9: Length frequencies of longfinned eels from stratum 1 of Mokoreta River.




Figure 10: Length frequencies of longfinned eels from stratum 1 of Waikaka Stream.




Figure 11: Length frequencies of longfinned eels from stratum 1 of Pomohaka River.












Figure 12: Length frequencies of longfinned eels from strata 1, 2, and 3 of Clutha River.



**Figure 13:** Length frequencies of longfinned eels from stratum 1 of Lake Wakatipu.



**Figure 14:** Length frequencies of longfinned eels from strata 3 and 4 of Taieri River.



**Figure 15:** Length frequencies of longfinned eels from strata 1, 2, and 3 of Waitaki River.




Figure 16: Length frequencies of longfinned eels from stratum 1 of Waimakariri River.





Figure 17: Length frequencies of longfinned eels from strata 2 and 3 of Grey River.






Figure 18: Length frequencies of longfinned eels from stratum 2 of Te Waihora.





Figure 19: Length frequencies of shortfinned eels from strata 1 and 2 of Waiau River.





Figure 20: Length frequencies of shortfinned eels from strata 1 and 2 of Aparima River.








Figure 21: Length frequencies of shortfinned eels from stratum 1 of Waituna Creek.




**Figure 22: Length frequencies of shortfinned eels from strata 1 and 2 of Oreti River.**




**Figure 23: Length frequencies of shortfinned eels from stratum 1 of Makarewa River.**






**Figure 24:** Length frequencies of shortfinned eels from stratum 1 of Mataura River.



**Figure 25:** Length frequencies of shortfinned eels from stratum 1 of Mokoreta River.



**Figure 26:** Length frequencies of shortfinned eels from stratum 1 of Waikaka Stream. Note the change of scale on the length axis.



**Figure 27: Length frequencies of shortfinned eels from strata 1, 2, and 3 of Clutha River.**




Figure 28: Length frequencies of shortfinned eels from stratum 1 of Pomahaka River.




Figure 29: Length frequencies of shortfinned eels from stratum 3 of Taieri River.




Figure 30: Length frequencies of shortfinned eels from stratum 1 of Grey River.






Figure 31: Length frequencies of shortfinned eels from strata 1, 2, and 3 of Waitaki River.








Figure 32: Length frequencies of shortfinned eels from strata 1, 2, and 3 of Te Waihora.





**Figure 33:** Length frequencies of shortfinned eels from stratum 4 of Te Waihora. Note the change of scale on the numbers axis.



**Figure 34:** Length frequencies of shortfinned eels from stratum 1 of Waimakariri River.

# Appendix 1: Gonad staging descriptions for longfinned and shortfinned eels

## Both species

### Immature or could not determined

These gonads can appear as a very thin, almost transparent, thread-like organ which cannot be differentiated into male or female for either species (usually where size is under 200 g). Commonly, gonads in this category were increasing in size with irregular segmentation but could not be clearly distinguished as male or female and were designated as stage 1 since gonads were of similar size to male and female stage 1. In 1996–97 eels in this category were staged from 1 to 3 where sex was not clear but gonads were clearly increasing in size relative to male and female staged eels (Beentjes & Chisnall 1998).

## Longfins

### Males

- |         |  |
|---------|--|
| Stage 1 | Regularly lobar Syrski organ developed even at very small sizes. Stage 1 organ is typically very narrow (2–3 mm), and often quite unrelated to fish length. Typified by flesh pink coloration. |
| Stage 2 | The lobar organ is usually slightly larger than in stage 1 (2–4 mm), but more dense in appearance, very obviously lobar and often ranges from dark pink to yellow.                             |
| Stage 3 | As for stage 2, but further developed in size and density.   |
| Stage 4 | Very large lobed organ, up to 12 mm or more in width, always dark red and with well developed vein network. Usually found in migrants and in longfins up to about 1300 g.                      |

### Notes on longfinned males

Most longfins less than about 350 g will be either stage 1 males or will be indeterminate as a result of having insufficiently lobed structure to the gonad strip. Regular lobe-type form is the key diagnostic feature, irrespective of coloration. Sometimes gonads in stage 2–3 sizes will appear male in density and form but will be incompletely lobed. These are assumed to be anomalous forms and are regarded as male. Some male fish have a number of external morphological features that assist in confirmation of sex, but must not be used as primary method of diagnosis: russet coloration, particularly about the underbelly; very tough belly skin in larger, more mature males; often a squared-off, or truncated looking tail; generally appear to have smaller relative head size and fatter bodies than females of the same length.

## **Females**

- Stage 1 Characterised by a thin (2–5 mm) ribbon, quite translucent and often broken in places, but not regularly segmented or lobar.
- Stage 2 Ribbon usually (but not necessarily) wider than stage 1 (about 4–10 mm), and displaying creamy pigmentation.
- Stage 3 Ribbon is wider than stage 2 (up to 20 mm) and is substantially folded or pleated. Completely opaque, cream or white.
- Stage 4 Very large, well developed white gonad with substantial vein network. Slightly granular texture in many stage 4 gonads, but not necessarily diagnostic. Usually found in migrants or near migrants.

## **Shortfins**

### **Males**

Shortfinned males are small, and therefore the development of the gonads is less conspicuous than in females, and most migrate before they recruit to the commercial fishery. Te Waihora is an exception since fishers are granted a dispensation to catch sublegal males from Ellesmere during the migration period of shortfinned males.

- Stage 1 Classic lobar organ of Syrski. Pale opaque lobes under 2 mm wide.
- Stage 2–4 Dark red, very lobar organs. Whether classed as stage 2, 3, or 4 depends on the relative size and depth of coloration. Nearly all males sampled were Te Waihora eels in the migrant morphological condition.

### **Females**

- Stage 1 Frilled ribbon 2–5 mm wide and translucent. Usually in small fish under 300g.
- Stage 2 Frilled ribbon 4–10 mm wide and pale, creamy colour. The most common stage in females.
- Stage 3 Frilled ribbon up to 15 mm wide. Usually with vein network. Completely opaque.
- Stage 4 Larger, heavy version of stage 3. Very white, and slightly granular structure. Usually observed only in migrants.



National Institute of  
Water and Atmospheric  
Research Limited